

FENOX EC

SPECIAL SENSITIVE AREA

USED AGAINST BED BUGS and DUST MITES
PUBLIC HYGIENE – ANIMAL HOUSING
BREEDING PREMISES – STORAGE PREMISES
DISPOSAL and STORAGE AREAS of WASTE

Formulation

Composition Etofenprox 300g/L / Natural pyrethrum 1g/L

Formulation Emulsifiable concentrate


Users Professionnals

Strenghts

WIDE SPECTRUM OF ACTIVITY: proven efficacy on flying (flies, mosquitoes...) and crawling insects (cockroaches, ants, fleas...), dust mites, bed bugs and also in stored foodstuffs (against Saw tooth beetle, Indian meal moth, Lesser grain borer, Rice weevil, Flour beetle, Cheese miteflour beetles...) and insects from animal housing areas and breeding premises (houseflies, litter beetles, red mites)

TREATMENTS IN VERY SENSITIVE AREA:

- Homes, offices, schools, hospitals and other public areas
- Packaged foodstuffs storage premises and food processing industry
- Breeding premises
- Disposal and storage areas of waste



*LD50= Lethal dose for 50% tested rats. The substance which has the highest LD50 is the less harmful

- Knock-down effet and long persistence of action (until 3 months) on porous and non-porous materials.
- Alternative in case of resistance to the classic pyrethrinoïds
- Etofenprox 300g/I + not classified ingredients = the power of the concentrate with a non agressive formulation.





References & Packaging

Ref. 18150: Bottle of 500mL

Ref. 18151: Water soluble sachets of 25mL x 10

Precautions of use

USE BIOCIDES SAFELY. Always read the label and product information before use. Read material data sheet. Use only as an insecticide.

Remove people, animals, aquariums and unpackaged foodstuffs from the area to be treated. Air the treated area before entering and staying. Wash hands and exposed skin with mild soap and water before eat, drink or smoke and when leaving work.

Do not apply directly on animals and unpackaged foodstuffs.

Do not expose to light, freezing, and heat. Keep out of the reach of children and animals.

Dispose of unused product and its containers in a safe manner, according to the national rules (in waste disposal center). In case of emergency, call the nearest poison center.





FENOX EC

SPECIAL SENSITIVE AREA

USED AGAINST BED BUGS and DUST MITES
PUBLIC HYGIENE – ANIMAL HOUSING
BREEDING PREMISES – STORAGE PREMISES
DISPOSAL and STORAGE AREAS of WASTE



Instructions for use

Surface spraying:

Spray the product after dilution in water (see table). Spray diluted FENOX EC on walls, floors, windows frames, carpets, balconies, terraces, and all other contaminated places, using a professional sprayer. Ventilate during the treatment (to avoid breathing liquid particles), then completely renew the air after treatment before reentering. Rinse the sprayer with water after use.

Volume and spatial treatment:

For an optimum efficacy, close all the openings such as doors, windows, gates... from the area before the treatment. The product can be used by cold fogging and thermo fogging. Use the product at the recommended dose (see table). Dilute in water if required, the quantity of water depends on the apparatus to be used and its flow. For instance, FENOX EC can be diluted at 25% when using a Swing Fog sprayer. Leave the treated premises after treatment and wait 4 to 6 hours. Thoroughly ventilate to renew completely the air, before re-entering (variable ventilation duration depending on level of ventilation and size of facility: 30mins to 2h). Rinse the apparatus with water after use. For a volume treatment, a complete disposable coverall, breathing mask, gloves and boots must be worn.

Application rates

| | SURFACE TREATMENT | VOLUME TREATMENT |
|---|------------------------------|---|
| PUBLIC HYGIENE | | |
| Mosquitoes | 50mL into 5L of water /100m² | 50mL / 1000m³ 30 to 60ml / 1ha |
| Other insects | 50mL into 5L of water /100m² | 200mL / 1000m ³ |
| STORAGE PREMISES, FOOD PROCESSING INDUSTRY, BAKERIES, SUPERMARKETS, | | |
| Storage premises | 50mL /100m² | / |
| Storage premises with packaged food in cardboard boxes | 50mL /100m² | 200mL / 1000m ³ 400mL/1000m ³ in case of infestation of cockroaches |
| BREEDING PREMISES | | |
| Animal housing | 50mL /100m ² | / |
| Equipment | 50mL /100m ² | / |
| Transport equipment | 50mL /100m ² | / |
| Food preparation areas | 50mL /100m ² | / |
| DISPOSAL AND STORAGE AREAS OF WASTE | | |
| Treatment areas | 50mL /100m ² | / |
| Transport equipment | 50mL /100m² | / |

